

Title: Mechanisms of nuclear positioning in skeletal muscle cells

Synopsis:

Position of the nucleus is important for cellular activities such as cell division, cell migration and multicellular organism development. The cytoskeleton plays an important role in nuclear positioning either by anchoring or moving the nucleus within the cell.

In our lab we are interested in understanding how does the cytoskeleton regulates nuclear positioning during muscle formation (1,2,3). We are also curious to know how mutations in proteins associated with muscular dystrophies interfere with nuclear position during myofiber formation (figure 1 and 2) (4).

We propose to study how these proteins are involved in the physical connection between the nucleus and the actin cytoskeleton during muscle formation. We use state-of-the-art microscopy techniques (multicolor time-lapse fluorescent microscopy, photo-activation and photoswitchable techniques, fluorescence recovery after photobleaching - FRAP) combined with molecular biology, biochemistry and micromanipulation (microinjection) approaches to address this process both *in vitro* and *in vivo* (figure 1 and 2). The mechanisms of nuclear positioning during muscle formation that we will identify are potential targets for therapies to inhibit abnormal nuclear positioning in muscle disorders.




Figure 1.

Isolated myofiber with peripheral nuclei.

Green - Gamma actin
Red – Microtubules
Blue - Nucleus




Figure 2- Skeletal muscle fiber, differentiated *in vitro*, with peripheral nuclei (blue) and fully formed triads (Red and Green).

Supervisor: Edgar Gomes, EGomes Lab, edgargomes@medicina.ulisboa.pt

Webpage of the group: <https://imm.medicina.ulisboa.pt/en/investigacao/labs/gomes-lab/>

Bibliography:

1. Gomes, E. R., Jani, S., and Gundersen, G. G. (2005). Nuclear movement regulated by Cdc42, MRCK, myosin, and actin flow establishes MTOC polarization in migrating cells. ***Cell*** 121, 451-63.
2. Luxton, G. W. G.*, Gomes, E. R.*, Folker, E. S., Vintinner, E., and Gundersen, G. G. (2010). Linear Arrays of Nuclear Envelope Proteins Harness Retrograde Actin Flow for Nuclear Movement. ***Science*** 329, 956-959. * co-first author
3. Metzger, T., Gache, V., Xu, M., Cadot, B., Folker, E., Richardson, B., Gomes, E.R.*, Baylies, M.K.*. MAP and Kinesin dependent nuclear positioning is required for skeletal muscle function. ***Nature***, 2012 *co-last and corresponding author
4. Borrego-Pinto, J., Jegou, T., Osorio, D., Aurade, F., Gorjánácz, M., Koch, B., Mattaj, I., and Gomes, E. R.. Samp1 is a new component of the TAN lines and is required for nuclear movement. ***Journal of Cell Science*** 2012
5. Falcone, S., Roman, W., Hnia, K., Gache, V., Didier, N., Lainé, J., Auradé, F., Marty, I., Nishino, I., Charlet-Berguerand, N., ER. Gomes (2014). N-WASP is required for Amphiphysin-2/BIN1-dependent nuclear positioning and triad organization in skeletal muscle and is involved in the pathophysiology of centronuclear myopathy. ***EMBO Mol Med*** 6, 1455–1475.

Remunerated or volunteer training:

Volunteer